

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

### STATE OF KNOWLEDGE – WHAT IS HAPPENING?

A very brief overview of the state of knowledge with respect to avian wildlife in the NWT is presented below. This overview is preliminary and not intended to be exhaustive.

→ **What are the baseline conditions with respect to avian wildlife?**

\* There are over 235 species of birds which migrate through or live in the NWT. The majority of these birds spend only the breeding season in the NWT. Increased mining, oil and gas exploration, as well as forestry and an expansion of road networks that always accompanies development, are significant concerns with regards to alteration or loss of habitat. Potential effects of climate change on habitat and distribution are also a concern.

\* The four main categories of avian wildlife— waterfowl, waterbirds, shorebirds and landbirds—are all found throughout the NWT. These categories are part of the classification used by the North American Bird Conservation Initiative. They are described below.

#### **Waterfowl and Waterbirds**

\* Waterfowl and waterbirds include gulls, terns, loons, ducks, geese, swans, cranes and rails. Generally they are abundant and widespread in the NWT. Baseline information is lacking for most species however some information is available through monitoring programs and surveys for specific species (mostly

### KEY MONITORING INDICATORS

<i>Population size and trend</i>	<i>Age structure</i>
<i>Number harvested</i>	<i>Habitat quality</i>
<i>Distribution and abundance</i>	<i>Food availability</i>
<i>Rate of breeding success</i>	<i>Presence of contaminants</i>
<b><i>Presence of diseases and parasites</i></b>	

in key breeding areas). Populations of some species of ducks (e.g. long tailed duck, lesser scaup, surf scoter, white-winged scoter, and northern pintail) have declined over the long term. Large-scale continental surveys are taking place. However, they cannot provide quality, local monitoring information. Two species of waterbirds may be at risk in the Northwest Territories: the yellow rail and American white pelican.

#### **Shorebirds**

\* Shorebirds include species such as sandpipers and plovers. They are widespread in the NWT, however baseline information is lacking and only general distribution can be described. Density estimates are available for a few locations only. Overall there appears to be a negative population trend in many shorebird species that are present in the NWT. An arctic shorebird monitoring program is now operational. It will determine current population levels of 17 species, and provide long-term trends in populations on the arctic breeding grounds. Testing is currently underway to develop a similar program in the boreal portion of the Northwest Territories. In the Northwest Territories the eskimo curlew is an endangered species (it has not been seen since the 1980s).

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

### **Landbirds**

\* Landbirds include songbirds, woodpeckers, cranes, owls and raptors. Some very preliminary information is available on population trends for song/forest birds, but it is not statistically rigorous. The endangered whooping crane (found only in Wood Buffalo National Park) is monitored and has a population size of 200 birds (2004 fall estimate). Two other species are designated as at/may be at risk: gray-headed chickadee and rusty blackbird.

\* Baseline information is available for some raptor species, such as the threatened *anatum* subspecies of peregrine falcon (81 nesting pairs were observed in 2000 throughout the Mackenzie River valley). Raptors tend to nest at historical nest sites from year to year. Monitoring raptors includes documenting nest site occupancy, reproduction rates (i.e. number of fledglings per nest) and production of fledglings within a geographic area. With the reduced use of organochlorine based pesticides in North America over the last 20 years, populations of raptors, including peregrine falcons, are stable or increasing and are considered to be healthy.

\* Species-specific or area specific research is being conducted as resources permit. Examples include work on rusty blackbird in 2006 and baseline data collection from near Fort Simpson, Norman Wells, and Colville Lake.

\* Additional data for landbirds are being collected as part of biological assessments of protected areas such as Edézhíe, Ts'ude'hilíne-Tuyetah, and others.

### → **What are the levels and trends of contaminants in avian wildlife?**

\* Baseline levels in contaminants in aquatic birds were evaluated in the late 1980s/early 1990s as part of a national contaminants survey. Contaminant levels in birds which are harvested were found to be low. However, other surveys of species higher in the food chain such as seabirds and gulls suggest that contaminant levels in some of these species are relatively high. Monitoring of contaminants in raptors in the NWT does not occur routinely therefore levels and trends are not available.

## RECENT AND CURRENT MONITORING

### **Ongoing monitoring programs with respect to avian wildlife in the NWT are found below.**

#### **Overall monitoring**

- ✓ Northwest Territories/Nunavut bird checklist survey (Canadian Wildlife Service since 1995)
  - This survey is volunteer based and serves to provide distribution and trend information for all bird species. Much of the data collected is historic (prior to 1995). This broad monitoring program will continue to be improved upon and expanded.
- ✓ North American breeding bird survey (Canadian Wildlife Service since 1966)
  - This survey is undertaken throughout North America mainly by volunteers using road based routes for data collection. The technique is reliable for some land birds

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

(mainly songbirds). There are five active routes in the NWT and others available for skilled observers.

- ✓ Christmas bird counts (Volunteer based since 1971).
  - Counts take place in Yellowknife, Hay River, Norman Wells, Fort Simpson and Fort Smith and provide data on winter bird populations.
- ✓ NWT species status rank infobase (Government of the Northwest Territories since 2000).
  - This infobase monitors the general status of species in the Northwest Territories. It is a significant source of information for assessing future monitoring of avian wildlife. The infobase is a searchable catalogue of information used to rank the status of species, thereby prioritizing them for further assessment and monitoring. The following biological indicators are used to rank species status: population size, number of occurrences, distribution, trend in population, trend in distribution, threats to population and threats to habitat. The general status ranks will be published every 5 years (see references in WGGNS 2006, methods in Carrière and Lange 2002).
- ✓ Wildlife effects monitoring program (BHP Diamonds Inc. since 1994).
  - The program monitors bird species, migrants or residents, found on the BHP claim block that may be affected by the EKATI diamond mine.
- ✓ Migratory bird and wildlife monitoring program for Fort Liard development project (Shiha Energy Transmission Ltd. since 2000).

- Birds and signs of other wildlife (tracks, pellet groups, etc.) are surveyed along the right-of-way. The right-of-way is treated as a disturbance feature, so information is collected in a way to show use of the feature, use of the edge, and use of the (undisturbed) surrounding habitat. This project was completed in 2004.

- ✓ Bird monitoring along the Chevron pipeline right-of-way, Fort Liard, Northwest Territories (Chevron Ltd. since 2000).
  - Point counts for bird species are conducted on the pipeline right-of-way and in adjacent habitat to compare use of the area by migratory birds. This project should be completed in 2005.

- ✓ Harvest study at Holman, Sachs Harbour, and Paulatuk (Canadian Wildlife Service and Wildlife Management Advisory Council) (since 1986)

### ***Aquatic birds***

- ✓ Duck banding projects (US Fish and Wildlife, Government of the Northwest Territories and Canadian Wildlife Service since 1995)
  - Duck banding occurs at Willow Lake, Mills Lake, and Stagg River, and is part of a continental duck banding program to monitor shifting distributions, survival (longevity) and harvest rates.
- ✓ Inventory of snow geese in western arctic (Canadian Wildlife Survey since 1976)
  - The inventory occurs at 5 year intervals in the Inuvialuit Settlement Region, but is of relevance to the NWT as birds migrate through and are harvested by residents.

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

### ✓ Monitoring Populations of waterfowl and other aquatic birds in the Inuvialuit Settlement

- The Mackenzie Delta, Tuktoyaktuk Peninsula, and neighbouring parts of the mainland Inuvialuit Settlement Region are one of the most important breeding areas for waterfowl and other aquatic birds in North America. Objectives of this long term monitoring study are to (1) document changes in numbers of several of species of waterfowl in one of the most important arctic breeding grounds; (2) evaluate the impacts of gas and oil development on migratory bird populations in the Inuvialuit Settlement Region; and (3) guarantee that sustainable and high numbers of waterfowl are maintained in the Mackenzie Delta region for the use of Inuvialuit and other Canadians.

### ✓ Evaluation effects of gas and oil development using tundra swans as an indicator species

- The Mackenzie Delta is one of the most important areas for nesting swans and other waterfowl in North America. The goal of this study is to determine the cumulative effects of oil/gas development and other environmental stressors on Tundra Swan populations and productivity, and the Mackenzie River delta ecosystem.

### ✓ Productivity of boreal forest duck and grebe populations (Canadian Wildlife Service since 1985)

- Long-term population trends and productivity of waterfowl in the boreal forest are being monitored along with factors which might limit population growth, habitat preferences and requirements of northern waterfowl, and methods for surveying northern aquatic birds.

### ✓ Annual spring aerial surveys of waterfowl breeding areas in North America (US Fish and Wildlife in cooperation with Canadian Wildlife Service since 1955)

- The NWT is included in this survey as an important breeding area.

### ✓ Five year surveys of Trumpeter Swans (Canadian Wildlife Service since 1985)

- Range wide surveys for swans are repeated at 5 year intervals to assess the population of this species.

### **Shorebirds**

### ✓ Arctic and boreal shorebird monitoring programs. (Since 2004. Canada-USA joint program).

### ✓ PRISM

- The Program for Regional and International Shorebird Monitoring (PRISM) is being carried out in NWT and Nunavut. Arctic PRISM is operational and uses a double-sampling methodology to estimate arctic-wide populations of 17 shorebird species. The entire arctic is sampled on a 10-year rotating basis; eventually trends in populations will emerge from this data. Information on habitat association and use is also being collected. Boreal PRISM is in the testing phase. Various survey methods will be tested in order to select the best methods for surveying the five priority species that breed in the boreal forest regions of Canada.

### **Land birds**

### ✓ Distribution and abundance of forest songbirds with relation to forest cover type in the Liard Valley (Canadian Wildlife Service 1998-2002, 2005, periodically resampled)

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

- The abundance and distribution of forest bird species in the Liard Valley was assessed as little was known about forest bird communities in this area. The potential for forest and petroleum development in the Liard Valley is high and such local studies provide baseline data for future reference.
- ✓ Ecology of whooping cranes – Annual surveys (Canadian Wildlife Service, Wood Buffalo National Park, and University of Alberta since the 1960s)
- ✓ Five year peregrine falcon surveys (Government of the Northwest Territories since the 1970s)
  - Peregrine falcon surveys (anatum and tundrius subspecies) are conducted every five years throughout the NWT. Surveys were conducted annually from the early 1970's to 1990 when a five year cycle was introduced. Historical peregrine nest sites are surveyed by air and ground to obtain information on occupancy rates, productivity and production of fledglings.
- ✓ Raptor monitoring at Daring Lake, Northwest Territories (Government of the Northwest Territories since 1995)
  - Raptors, including peregrine falcons, gyr falcons and rough legged hawks, are monitored annually in the Daring Lake area. A productivity survey is conducted by helicopter in July to document occupancy, productivity and production of young birds.
- ✓ Raptor monitoring (Diavik Diamond Mines Inc. since 2000)
  - Annual surveys of peregrine falcons began to determine the effects of mine development on nesting peregrines.

### GAPS AND RECOMMENDATIONS FOR MONITORING

**A list of monitoring gaps and recommendations for future monitoring under the NWT Cumulative Impact Monitoring Program is found below.**

#### **Gaps**

- Large gaps in monitoring of distribution, abundance, population size, breeding success and condition/health for most bird species; most information is site specific and many studies cannot be combined to obtain any regional population trends.
- Significant gaps for waterfowl and waterbirds which have been declining such as scaups, scoters and long tailed ducks
- Many monitoring programs are carried out sporadically from year to year due to lack of consistent core funding
- Too few breeding bird survey routes for reliable population trend analyses
- Abundance, distribution and population size of raptors is unknown in many regions which have never been surveyed

#### **Recommendations**

- Species specific long term monitoring of distribution, abundance, population size and breeding success, particular for declining species

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

- Monitoring changes in bird communities and their habitat in relation to possible climate-induced changes
- Evaluation of baseline contaminant levels in aquatic bird species, to compare them to results from late 1980s/ early 1990s
- A community-based seasonal monitoring program to document bird phenology (arrival, departure, nesting dates), distribution (presence/absence) and breeding status, for both game and nongame species in a similar format to harvest studies
- A community-based program to monitor qualitative changes in bird condition, including body fat indices, parasites and diseases (mainly for game species – similar to harvest studies)

### REFERENCES

Relevant monitoring reports, past monitoring programs, research documents, and scientific publications are found below.

#### OVERALL

Working Group on General Status of NWT Species. 2006. **NWT Species 2006-2010 – General Status Ranks of Wild Species in the Northwest Territories**. Dept. of Environment and Natural Resources, Government of the NWT, Yellowknife, NT.

Canadian Wildlife Service 2006. **Shorebird program strategy and action plan**. Northern Conservation Division, Canadian Wildlife Service, Yellowknife, NT.

Canadian Wildlife Service 2002. **Landbird program strategy and action plan**. Northern Conservation Division, Canadian Wildlife Service, Yellowknife, NT.

Golder Associates Ltd. (2001). **Bird monitoring along the Chevron pipeline right-of-way, Fort Liard, Northwest Territories**. Calgary, AB. 8 pp. (see Current Monitoring for description)

Government of the Northwest Territories (2000). **NWT Species 2000: General status ranks of wild species in the Northwest Territories**. (see Current Monitoring for description)

Hornbeck, G.E. and D. Soprovich (2004). **Migratory bird monitoring program for the Fort Liard development project: Final report year five**. Prepared for Shiha Energy Transmission, Ltd., Calgary, by Wildlife and Company Ltd in association with Bluestem Wildlife Services. 33 pp.+ appendices (see Current Monitoring for description)

Working Group on General Status of NWT Species (WGGSNS) 2006. **NWT Species 2006-2010 – General Status of Wild Species in the Northwest Territories**, Department of Environment and Natural Resources, GNWT, Yellowknife, NT. 111 pp.

#### AQUATIC BIRDS

Braune, B.M., Hobson, K.A. and B.J. Malone (2005). **Regional differences in collagen stable isotope and trace element profiles in populations of long-tailed duck breeding in the Canadian Arctic**. *Science of the Total Environment* 346: 156-168.

Braune, B.M. and B.J. Malone (2006). **Mercury and selenium in livers of waterfowl harvested in northern Canada**. *Archives of Environmental Contamination and Toxicology* 50: 284-289.

Braune, B.M. and B.J. Malone (2006). **Organochlorines and mercury in waterfowl harvested in Canada**. *Environmental Monitoring and Assessment* 114: 331-359.

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

Braune, B.M. and B.J. Malone (2006). **Organochlorines and trace elements in upland game birds harvested in Canada.** *Science of the Total Environment* 363: 60-69.

Bromley, R.G. (1993). **Comparison of results from two waterfowl harvest studies in three Inuvialuit communities, Northwest Territories, 1988-1990.** Government of the Northwest Territories. 20pp.

Canadian Wildlife Service. **Spring migration of waterfowl in the Yellowknife-Thor Lake area, Northwest Territories (1986-1988).**

Canadian Wildlife Service. **Assessment of Mills Lake as a staging area for migrating waterfowl and shorebirds (1994-1997).**

Canadian Wildlife Service (1998). **Status of waterfowl in the Inuvialuit Settlement Region.** Unpublished report. Canadian Wildlife Service, Yellowknife and Inuvialuit Wildlife Management Advisory Council, Inuvik, NT.

Canadian Wildlife Service Waterfowl Committee (2000). **Population Status of Migratory Game Birds in Canada: November 2000.** Canadian Wildlife Survey Migratory Birds Regulatory Report Number 1. Ottawa, ON.  
<http://www.cws-scf.ec.gc.ca/canbird/status/nov00/nov2000p.pdf>

Canadian Wildlife Service. **Harvest of king and common eiders at Holman, NWT (1996-1999)**

Ducks Unlimited. **Distribution, abundance and nesting success of waterfowl in the Bracket Lake wetland complex (1997-1999).**

Fournier, M.A. and R.G. Bromley (1996). **Status of the Harlequin Duck, *Histrionicus histrionicus*, in the western Northwest Territories.** *The Canadian Field-Naturalist* 110(4): 638-641.

Fournier, M.A. and J.E. Hines (1996). **Changed status of the Hooded Merganser, *Lophodytes cucullatus*, in the Yellowknife**

**area, Northwest Territories.** *The Canadian Field-Naturalist* 110(4): 713-714.

Fournier, M.A. and J.E. Hines (1998). **Productivity and population increase of subarctic breeding canvasbacks.** *Journal of Wildlife Management* 62(1): 179-184.

Fournier, M.A. and J.E. Hines (1998). **Breeding ecology and status of the Red-necked Grebe, *Podiceps grisegena*, in the subarctic of the Northwest Territories.** *Canadian Field-Naturalist* 112(3): 474-480.

Fournier, M.A. and J.E. Hines (1999). **Breeding ecology of the Horned Grebe (*Podiceps auritus*) in subarctic wetlands.** Occasional Paper Number 99, Canadian Wildlife Service, Ottawa. 32 pp.

Fournier, M.A. and J.E. Hines. 2001. **Breeding ecology of sympatric Greater and Lesser Scaup (*Aythya marila* and *Aythya affinis*) in the subarctic Northwest Territories.** *Arctic* 54(4):444-456.

Fournier, M.A. and J.E. Hines (in prep.). **Waterfowl studies on the North Arm of Great Slave Lake 1989-1995.** Occasional Paper, Canadian Wildlife Service, Ottawa, ON.

Hines, J.E. and M.O. Wiebe Robertson (editors.), 2006. **Surveys of geese and swans in the Inuvialuit Settlement Region, Western Canadian Arctic, 1989-2001.** *Can. Wildl. Serv. Occas. Paper No. 112.* *Can. Wildl. Serv.*, Ottawa. 73 pp.

Kerbes, R.H., K.M. Meeres, and J.E. Hines (editors). 1999. **Distribution, survival, and numbers of Lesser Snow Geese of the Western Canadian Arctic and Wrangel Island, Russia.** Canadian Wildlife Service Occasional Paper #98. Ottawa, Ontario.

Wayland, M., K.A. Hobson, and J. Sirois (2000). **Environmental contaminants in colonial waterbirds from Great Slave Lake, NWT:**

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

*Spatial, temporal and food-chain considerations. Arctic 53(3): 221-233.*

### SHOREBIRDS

Bart, J., V. Johnston and P. Smith. 2005. **Design of a long-term survey for shorebirds in the arctic.** Unpubl. Manuscript, available from Canadian Wildlife Service, Yellowknife.

Canadian Wildlife Service. **Lesser yellowlegs monitoring pilot project.** Conducted in 2000 near Yellowknife, NT. Unpubl. Manuscript available from CWS Yellowknife.

Sinclair, P., Y. Aubry, J. Bart, V. Johnston, R. Lanctot, B. McCaffrey, K. Ross, P. Smith and L. Tibbitts, 2004. **Boreal shorebirds – An Assessment of Conservation Status and Potential for Population Monitoring.** Unpubl. Manuscript, available from Canadian Wildlife Service, Yellowknife.

**Population estimates of shorebirds in the Mackenzie Delta, 2005-2006.** Unpubl. Ms available from CWS Yellowknife.

PRISM database. **Database of shorebird habitats, densities, and population estimates collected in various parts of the NWT and Nunavut, 2001-2006.** CWS Yellowknife.

### LAND BIRDS

Bromley, R.G. and S.B. Mathews (1988). **Status of peregrine falcons in the Mackenzie River Valley, NWT, 1969-1985.** In: Cade, T. *Peregrine Falcon Populations: Their Management and Recovery.* The Peregrine Fund, Boise, Idaho, U.S.A

Cooper, John M.; Wheatley, Matthew; Chytky, Paul A.; Deans, Aaron; Holschuh, Carmen ; and Beauchesne, Suzanne M. 2004. **Potential impacts on birds in the Mackenzie Valley from Pipeline Clearings for the Mackenzie Gas Project.** Technical Report Series No. 442. Canadian Wildlife Services, Prairie and Northern Region, Yellowknife.

Government of the Northwest Territories. **Monitoring nesting success of Mackenzie Valley peregrine falcons (1989-1995).**

Government of the Yukon-Renewable Resources. **Gyrfalcon population monitoring, North Slope (1972-1992).**

Holroyd, G.L. and U. Banasch (1996). **The 1990 Canadian peregrine falcon (*Falco peregrinus*) survey.** *Journal of Raptor Research* 30(3): 145-156.

Machtans, C.S. and P.B. Latour. 2003. **Boreal Forest Songbird Communities of the Liard Valley, Northwest Territories.** *The Condor* 105(1):27-44.

Machtans, C.S. and D.A. Kirk 2001. **Can we use songbirds to monitor and assess cumulative effects of development in the boreal forest?** Internal report, Canadian Wildlife Service, Yellowknife, NT.

Machtans, C.S., S.L. Van Wilgenburg, L.A. Armer, and K.A. Hobson. 2007. **Retrospective comparison of the occurrence and abundance of Rusty Blackbird in the Mackenzie Valley, Northwest Territories.** *Avian Conservation and Ecology* (in press, February 2007).

Machtans, C.S. 2006. **Songbird response to seismic lines in the western boreal forest: A manipulative experiment.** *Canadian Journal of Zoology* 84: 1421-1430.

Savignac, C. and C.S. Machtans. 2006. **Habitat requirements of Yellow-bellied Sapsucker, *Sphyrapicus varius*, in boreal mixedwood forests of northwestern Canada.** *Canadian Journal of Zoology* 84: 1230-1239.

Machtans, C. S. (2000). **Extra-limital observations of broad-winged hawk, *Buteo platypterus*, Connecticut warbler, *Oporornis agilis*, and other bird observations from the Liard**

## Valued Component – Other Wildlife (Avian) (FINAL DRAFT)

**Valley, Northwest Territories.** *Canadian Field-Naturalist* 114(4): 359-367.

Matthews, S.B. (1989). **Norman Wells pipeline project raptor monitoring program, 1980 to 1988.** Government of the Northwest Territories. 46pp.

Matthews, S.B. and S. Carriere (2000). **2000 Mackenzie Valley, NWT – Peregrine falcon survey.** Government of the Northwest Territories.

Matthews, S.B. and S. Carriere (2005). **2005 Mackenzie Valley, NWT – Peregrine falcon survey.** Government of the Northwest Territories.

Wakelyn, L.A., C.C. Shank, B.T. Elkin and D.C. Dragon (1999). **Contaminant levels in willow ptarmigan (*Lagopus lagopus*) from the western Canadian Arctic.** *Canadian Field-Naturalist* 113 (2): 215-220.